

Published in final edited form as:

*Nat Genet.* 2007 October ; 39(10): 1256–1260.

## Diet and the evolution of human amylase gene copy number variation

George H. Perry<sup>1,2,\*</sup>, Nathaniel J. Dominy<sup>3,\*</sup>, Katrina G. Claw<sup>1,4</sup>, Arthur S. Lee<sup>2</sup>, Heike Fiegler<sup>5</sup>, Richard Redon<sup>5</sup>, John Werner<sup>4</sup>, Fernando A. Villanea<sup>3</sup>, Joanna L. Mountain<sup>6</sup>, Rajeev Misra<sup>4</sup>, Nigel P. Carter<sup>5</sup>, Charles Lee<sup>2,7,†</sup>, and Anne C. Stone<sup>1,†</sup>

<sup>1</sup>*School of Human Evolution and Social Change, Arizona State University, Tempe, AZ 85287, USA*

<sup>2</sup>*Department of Pathology, Brigham and Women's Hospital, Boston, MA 02115, USA*

<sup>3</sup>*Department of Anthropology, University of California, Santa Cruz, CA 95064, USA*

<sup>4</sup>*School of Life Sciences, Arizona State University, Tempe, AZ 85287, USA*

<sup>5</sup>*The Wellcome Trust Sanger Institute, The Wellcome Trust Genome Campus, Hinxton, Cambridge, CB10 1SA, United Kingdom*

<sup>6</sup>*Department of Anthropological Sciences, Stanford University, Stanford, CA 94305, USA*

<sup>7</sup>*Harvard Medical School, Boston, MA 02115, USA*

### Abstract

Starch consumption is a prominent characteristic of agricultural societies and hunter-gatherers in arid environments. In contrast, rainforest and circum-arctic hunter-gatherers and some pastoralists consume much less starch<sup>1-3</sup>. This behavioral variation raises the possibility that different selective pressures have acted on amylase, the enzyme responsible for starch hydrolysis<sup>4</sup>. We found that salivary amylase gene (*AMY1*) copy number is correlated positively with salivary amylase protein levels, and that individuals from **populations with high-starch diets have on average more *AMY1* copies than those with traditionally low-starch diets**. Comparisons with other loci in a subset of these populations suggest that the level of *AMY1* copy number differentiation is unusual. **This example of positive selection on a copy number variable gene is one of the first in the human genome.** **Higher *AMY1* copy numbers and protein levels likely improve the digestion of starchy foods**, and may buffer against the fitness-reducing effects of intestinal disease.

Hominin evolution is characterized by significant dietary shifts, facilitated in part by the development of stone tool technology, the control of fire, and most recently the domestication of plants and animals<sup>5-7</sup>. Starch, for instance, has become an increasingly prominent component of the human diet, particularly among agricultural societies<sup>8</sup>. It stands to reason, therefore, that studies of the evolution of amylase in humans and our close primate relatives may provide insight into our ecological history. Because the human salivary amylase gene (*AMY1*) shows extensive variation in copy number<sup>9,10</sup>, we first assess whether a functional relationship exists between *AMY1* copy number and the level of amylase protein expression in

**Corresponding Author:** Dr. Nathaniel J. Dominy, Department of Anthropology, University of California, Santa Cruz, CA 95064, Telephone: (831) 459-2541, Fax: (831) 459-5900, E-mail: njdominy@ucsc.edu.

\*G.H.P. and N.J.D. contributed equally to this work

†C.L. and A.C.S. contributed equally to this work

**Author Contributions:** G.H.P., N.J.D., C.L. and A.C.S. designed the study, G.H.P., F.A.V., J.L.M. and A.C.S. collected the samples, G.H.P. and A.S.L. performed qPCR experiments, J.W. performed western blot experiments, G.H.P. performed fiber FISH experiments, H.F. and R.R. performed and analyzed aCGH experiments, K.G.C. performed nucleotide sequencing experiments, G.H.P. performed data analyses, R.M., N.P.C., C.L. and A.C.S. supervised the experiments and analyses, and G.H.P. and N.J.D. wrote the paper.

saliva. We then determine if *AMY1* copy number differs among modern human populations with contrasting levels of dietary starch.

We estimated diploid *AMY1* gene copy number for 50 European-Americans using an *AMY1*-specific real-time quantitative polymerase chain reaction (qPCR) assay. We observed extensive variation in *AMY1* copy number in this population sample (Fig. 1a and Supplementary Table 1 online), consistent with previous studies<sup>10,11</sup>. Next, we performed western blot experiments with saliva samples from the same individuals in order to estimate salivary amylase protein levels (Fig. 1b). These experiments revealed a significant positive correlation between salivary amylase gene copy number and protein expression level ( $P < 0.001$ ; Fig. 1c).

While there is a considerable range of variation in dietary starch intake among human populations, a distinction can be made between “high-starch” populations for which starchy food resources comprise a substantial portion of the diet, and the small fraction of “low-starch” populations with traditional diets that incorporate relatively few starchy foods. Such diets instead emphasize proteinaceous resources (e.g., meats and blood) and simple saccharides (e.g., from fruit, honey and milk). To determine if *AMY1* copy number differs among populations with high- and low-starch diets, we estimated *AMY1* copy number in three high-starch and four low-starch population samples. Our high-starch sample included two agricultural populations, European-Americans ( $n = 50$ ) and Japanese ( $n = 45$ ), and Hadza hunter-gatherers who rely extensively on starch-rich roots and tubers ( $n = 38$ )<sup>12</sup>. Low-starch populations included Biaka ( $n = 36$ ) and Mbuti ( $n = 15$ ) rainforest hunter-gatherers, Datog pastoralists ( $n = 17$ ), and the Yakut, a pastoralist/fishing society ( $n = 25$ ). Additional details on the diets of these populations are provided in Supplementary Table 2 online. We found that mean diploid *AMY1* copy number is greater in high-starch populations (Fig. 2 and Supplementary Fig. 1 online). Strikingly, the proportion of individuals from the combined high-starch sample with at least 6 *AMY1* copies (70%) is nearly 2 times greater than that for low-starch populations (37%). To visualize the allele-specific number and orientation of *AMY1* gene copies, we performed high-resolution fluorescence *in situ* hybridization on stretched DNA fibers (fiber FISH); these results were consistent with diploid *AMY1* copy number estimates from our qPCR experiments (Fig. 3a,b).

The among-population patterns of *AMY1* copy number variation do not fit expectations under a simple regional-based model of genetic drift: our high- and low-starch samples include both African and Asian populations, suggesting that diet more strongly predicts *AMY1* copy number than geographic proximity. Based on this observation, we hypothesized that natural selection may have influenced *AMY1* copy number in certain human populations. However, we cannot rigorously test such a hypothesis on the basis of our qPCR results alone, in part because we lack comparative data from other loci. Therefore, we next performed array-based comparative genomic hybridization (aCGH) on the Yakut population sample with a Whole Genome TilePath (WGTP) array platform that was previously used by Redon and colleagues<sup>11</sup> to describe genome-wide patterns of copy number variation in 270 individuals (the HapMap collection), including the same Japanese population sample as in our study. For the Yakut aCGH experiments, we used the same reference DNA sample (NA10851) as in the previous study<sup>11</sup>, facilitating comparisons of Japanese and Yakut relative intensity  $\log_2$  ratios for the 26,574 bacterial artificial chromosome (BAC) clones on the array, including two clones mapped to the *AMY1* locus.

Results from the two *AMY1*-mapped clones on the WGTP array supported our original observations: the  $\log_2$  ratios were strongly correlated with the qPCR estimates of *AMY1* diploid copy number (Supplementary Fig. 1 online), and the population mean  $\log_2$  ratios for both clones were greater for the Japanese sample (Fig. 4a and Supplementary Fig. 1 online). More importantly, with the WGTP data we were able to compare the level of population differentiation at the *AMY1* locus to other loci in the genome for the two Asian population

samples in our study. We would expect the magnitude and direction of the Japanese-Yakut mean log<sub>2</sub> ratio difference for the *AMY1*-mapped clones to be similar to those for other copy number variable clones, if these CNVs have experienced similar evolutionary pressures. However, the two *AMY1*-mapped clones are significant outliers in this distribution (Fig. 4b and Supplementary Fig. 2 online), leading us to reject this null hypothesis. In addition, we considered a database of genotypes for 783 genome-wide microsatellites for the same Yakut individuals and a different Japanese population sample<sup>13</sup>, because microsatellite loci are usually multi-allelic (as is the *AMY1* locus). We found that the level of Japanese-Yakut differentiation at the *AMY1* locus exceeds that for >97% of the microsatellite loci (Supplementary Fig. 3 online). Although this result should be interpreted with caution because we do not know whether *AMY1* copy number and microsatellite mutation rates and patterns are similar, this finding is consistent with our results from the genome-wide WGTP comparison.

These observations suggest that natural selection has shaped *AMY1* copy number variation in either the Japanese or the Yakut, or in both populations. We cannot fully test the null hypothesis for the other high- and low-starch populations in our study, but the patterns of copy number variation we observed in these populations are similar to those for the Japanese and Yakut and therefore may also reflect non-neutral evolution. We favor a model in which *AMY1* copy number has been subject to positive or directional selection in at least some high-starch populations but has evolved neutrally (i.e., through genetic drift) in low-starch populations. Although it is possible that lower *AMY1* gene copy numbers have been favored by selection in low-starch populations, such an interpretation is less plausible for the simple reason that excessive amylase production is unlikely to have a significant negative effect on fitness. Furthermore, several lines of evidence offer mechanisms by which higher salivary amylase protein levels may confer a fitness advantage for individuals with a high-starch diet. First, a significant amount of starch digestion occurs in the mouth during mastication<sup>14</sup>. For example, blood glucose levels have been shown to be significantly higher when high-starch foods such as corn, rice, and potatoes (but not apples) are first chewed and then swallowed, rather than swallowed directly<sup>15</sup>. In addition, it has been suggested that oral digestion of starch is critically important for energy absorption during episodes of diarrhea<sup>4</sup>. Diarrheal diseases can have a significant effect on fitness; for example, such diseases caused 15% of worldwide deaths among children younger than 5 years as recently as 2001<sup>16</sup>. Lastly, salivary amylase persists in the stomach and intestines after swallowing<sup>17</sup>, thereby augmenting the enzymatic activity of pancreatic amylase in the small intestine. Higher *AMY1* copy number and a concomitant increase in salivary amylase protein level are therefore likely to improve the efficiency with which high-starch foods are digested in the mouth, stomach, and intestines, and may also buffer against the potential fitness-reducing effects of intestinal disease.

To understand better the evolutionary context of human *AMY1* copy number variation, we analyzed patterns of *AMY1* copy number variation in chimpanzees (*Pan troglodytes*) and bonobos (*Pan paniscus*). In contrast to the extensive copy number variation we observed in humans, each of 15 wild-born western chimpanzees (*P. t. verus*) showed evidence of only 2 diploid *AMY1* copies (Fig. 3c and Supplementary Fig. 4 online), which is consistent with previous findings<sup>18-21</sup>. Although we observed evidence of a gain in *AMY1* copy number in bonobos relative to chimpanzees (Supplementary Fig. 4 online), our sequence-based analyses suggest that each of these *AMY1* copies has a disrupted coding sequence and may be non-functional (Supplementary Fig. 5 online). Therefore, the average human has ~3 times more *AMY1* copies than chimpanzees, and bonobos may not have salivary amylase at all. Outgroup comparisons with other great apes suggest that *AMY1* copy number was most likely gained in the human lineage, rather than lost in chimpanzees<sup>21,22</sup>. Given that *AMY1* copy number is positively correlated with salivary amylase protein level in humans, it stands to reason that the human-specific increase in copy number may explain, at least in part, why salivary amylase

protein levels are ~6-8 times higher in humans than in chimpanzees<sup>23</sup>. These patterns are consistent with the general dietary characteristics of *Pan* and *Homo*; chimpanzees and bonobos are predominantly frugivorous and ingest little starch relative to most human populations<sup>24</sup>. Considering other primates, while New World monkeys do not produce salivary amylase and tend to consume little starch, cercopithecines (a subfamily of Old World monkeys including macaques and mangabeys) have relatively high salivary amylase expression, even compared to humans<sup>23</sup>. Although the genetic mechanisms are unknown, this expression pattern may have evolved to facilitate the digestion of starchy foods (such as the seeds of unripe fruits) stowed in the cheek pouch, a trait that among primates is unique to cercopithecines<sup>25</sup>.

The initial human-specific increase in *AMY1* copy number may have been coincident with a dietary shift early in hominin evolutionary history. For example, it is hypothesized that starch-rich plant underground storage organs (USOs) were a critical food resource for early hominins<sup>26,27</sup>. Changes in USO consumption may even have facilitated the initial emergence and spread of *Homo erectus* out of Africa<sup>5,28</sup>. Yet such arguments are difficult to test, mainly because direct evidence for the use of USOs is difficult to obtain, particularly for more remote time periods. USOs themselves are perishable, as are many of the tools used to collect and process them. Therefore, understanding the timing and nature of the initial human-lineage *AMY1* duplications may provide insight into our ecological and evolutionary history. The low level of nucleotide sequence divergence among the three *AMY1* gene copies found in the human genome reference sequence (hg18;  $d = 0.00011$  to  $0.00056$ ) implies a relatively recent origin that may be within the timeframe of modern human origins (i.e., within the last ~200,000 years; based on human-chimpanzee *AMY1*  $d = 0.027$  and a 6 MYA estimate for divergence of the human and chimpanzee lineages). However, given the possibility for gene conversion, we do not necessarily consider this estimate to be reliable. The generation of *AMY1* sequences from multiple human individuals may ultimately help to shed light on this issue.

In summary, we have shown that the pattern of variation in copy number of the human *AMY1* gene is consistent with a history of diet-related selection pressures, demonstrating the importance of starchy foods in human evolution. While the amylase locus is one of the most variable in the human genome with regard to copy number<sup>10</sup>, it is by no means unique; a recent genome-wide survey identified 1,447 copy number variable regions among 270 phenotypically normal human individuals<sup>11</sup>, and many more such regions will likely be discovered with advances in copy number variation detection technology. It is reasonable to speculate that copy number variants other than *AMY1* are or have been subject to strong pressures of natural selection, particularly given their potential influence on transcriptional and translational levels (e.g., ref. <sup>29</sup>). The characterization of copy number variation among humans and between humans and other primates promises to offer considerable insight into our evolutionary history.

## Methods

### Samples

Buccal swabs and saliva were collected under informed consent from 50 European-Americans age 18-30 (Arizona State University IRB protocol no. 0503002355). Saliva was collected for 3 min from under the tongue. Buccal swabs were collected from the Hadza ( $n = 38$ ) and Datog ( $n = 17$ ) from Tanzania (Stanford University IRB protocol no. 9798-414). Genomic DNAs from Biaka (Central African Republic;  $n = 32$ ), Mbuti (Democratic Republic of Congo;  $n = 15$ ) and Yakut (Siberia;  $n = 25$ ) are from the HGDP-CEPH Human Genome Diversity Cell Line Panel. Lymphoblastoid cell lines from 45 Japanese, 4 additional Biaka, and the donor for the chimpanzee genome sequence (Clint) were obtained from the Coriell Institute for Medical Research. Whole bloods were collected during routine veterinary examinations from chimpanzees and bonobos housed at various zoological and research facilities. Two additional

bonobo samples were obtained from the Integrated Primate Biomaterials and Information Resource. DNA was isolated using standard methods.

### Copy number estimation

Primers for qPCR (Supplementary Table 3 online) were designed to be specific to *AMY1* (i.e., sequence mismatches with *AMY2A* and *AMY2B*) based on the human and chimpanzee reference genome sequences. A previous study reported a single (haploid) copy of *AMY1* for one chimpanzee<sup>18</sup>, and a recent analysis by Cheng et al.<sup>19</sup> found no evidence of recent *AMY1* duplication for Clint. We used fiber FISH to confirm that Clint has two diploid copies of *AMY1* (Fig. 3c). Therefore, we were able to estimate diploid copy number based on relative *AMY1* quantity for human DNAs compared to a standard curve constructed from the DNA of Clint. A fragment from the *TP53* gene was also amplified to adjust for DNA dilution quantity variation. Samples were run in triplicate and standards in duplicate. Experiments were performed and analyzed as described<sup>20</sup>.

### Western blot analysis

Protein samples were prepared by solubilizing saliva samples in 2% sodium dodecyl sulfate (SDS) and heating at 100°C for 5 min. These samples were analyzed on mini SDS-polyacrylamide gels and transferred to polyvinylidene difluoride (PVDF) membranes (Immobilon-Millipore). For quantification purposes, a human salivary amylase protein sample of known quantity (Sigma) was run on each gel, with 5 µL of saliva for each sample. After transfer, the membranes were incubated for 1.5 hours with primary antibodies raised against human salivary amylase (Sigma). The membranes were washed and goat anti-rabbit alkaline phosphatase conjugated IgG secondary antibodies (Pierce) were added for 1 hour. The membranes were exposed to ECF substrate (Amersham Biosciences) for 5 min and analyzed using a phosphorimager. Quantification of protein bands was performed using ImageQuant software (Molecular Dynamics).

### Fiber FISH

DNA fibers were prepared by gently lysing cultured lymphoblast cells with 300 µl Cell Lysis Buffer (Gentra Systems) per 5 million cells. 10 µl of lysate was placed on a poly-L-lysine coated slide (LabScientific) and mechanically stretched with the edge of a coverslip. After 30 sec, 300 µl of 100% methanol was applied to fix the fibers. Slides were dried at 37° C for 5 min and then stored at room temperature (RT).

PCR product probes were made from (i) the entire *AMY1* gene itself (~10 kb; red in images), and (ii) the retrotransposon found directly upstream of all *AMY1* copies but not pancreatic amylase genes or amylase pseudogenes (~8 kb; green in images); while the gene probe may not be specific to *AMY1* under all hybridization conditions (*AMY1* sequence divergence with *AMY2A* and *AMY2B* = 7.5% and 7.1%, respectively), the upstream probe is. We used long-range followed by nested PCR for each region (primers and conditions are provided in Supplementary Table 3 online). PCR products were purified with DNA Clean and Concentrator columns (Zymo).

For each nested PCR product, 750 ng was combined with 20 µl 2.5x random primer (BioPrime aCGH Labeling Module, Invitrogen) in 39 µl total volume, placed at 100° C for 5 min, and then ice for 5 min. Next, 5 µl 10x dUTP and 1 µl Exo-Klenow Fragment (BioPrime Module), and either 5 µl (5 nmol) Biotin-16-dUTP (Roche; gene probe) or 5 µl (5 nmol) Digoxigenin-11-dUTP (Roche; upstream probe) were added, and incubated at 37° C for 5 hours. Labeled products were purified with Microcon Centrifugal Filter Devices (Millipore) using 3 washes of 300 µl 0.1x SSC, eluted with 50 µl H<sub>2</sub>O. For each 1 µg of labeled DNA, we added 10 µg human Cot-1 DNA (Invitrogen).

For each experiment, 500 ng of labeled DNA from each of the nested PCR reactions were combined, lyophilized, reconstituted in 10 µl hybridization buffer (50% formamide, 20% dextran sulfate, 2x SSC), and added to the slide (18×18 mm cover glass; Fisher). Fibers and probes were co-denatured (95° C for 3 min) and hybridized in a humidified chamber (37° C for 40 hours). The slide was washed in 0.5x SSC at 75° C for 5 min followed by 3 washes in 1x PBS at RT for 2 min each. Next, fibers were incubated with 200 µl CAS Block (Zymed) and 10% Normal Goat Serum (Zymed) for 20 min at RT under a HybriSlip (Invitrogen). We used a 3-step detection/amplification (with reagents in 200 µl CAS Block/ 10% Normal Goat Serum). Each step was 30 min at RT under a HybriSlip followed by 3 washes in 1x PBS for 2 min each at RT: (i) 1:500 Anti-digoxigenin-fluorescein, Fab fragments (Roche) and 1:500 Streptavidin, Alexa Fluor 594 conjugate (Invitrogen); (ii) 1:250 Rabbit anti-FITC antibody (Zymed) and 1:500 Biotinylated anti-streptavidin (Vector Laboratories); (iii) 1:100 Goat anti-rabbit IgG-FITC (Zymed) and 1:500 Streptavidin, Alexa Fluor 594 conjugate. Images were captured on an Olympus BX51 fluorescent microscope with an Applied Imaging camera and analyzed with Applied Imaging's Genus software.

### aCGH analysis

For aCGH experiments we used a large-insert clone DNA microarray covering the human genome in tiling path resolution<sup>30</sup>. Test (Yakut individuals) and reference (NA10851) genomic DNA samples were labeled with Cy3-dCTP and Cy5-dCTP, respectively (NEN Life Science Products) and co-hybridized to the array. For each sample, a duplicate experiment was performed in dye-swap to reduce false-positive error rates. Labeling, hybridization, washes, and analyses were performed as described<sup>11,30</sup>.

### Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


### Acknowledgments

We are grateful to all our study participants. We thank Drs. H. Cann and C. de Toma of the Fondation Jean Dausset (CEPH), the Cincinnati Zoo, Lincoln Park Zoo, New Iberia Research Center, Primate Foundation of Arizona, Southwest Foundation for Biomedical Research, Coriell Institute for Medical Research, and the Integrated Primate Biomaterials and Information Resource for samples. C. Tyler-Smith and Y. Gilad provided helpful comments on a previous version of the manuscript. We would also like to thank the Wellcome Trust Sanger Institute Microarray Facility for printing the arrays and T. Fitzgerald and D. Rajan for technical support. This study was funded by grants from the L.S.B. Leakey Foundation and Wenner-Gren Foundation (to N.J.D.), the Department of Pathology, Brigham & Women's Hospital (to C.L.), the National Institutes of Health (to the University of Louisiana at Lafayette New Iberia Research Center; nos. RR015087, RR014491 and RR016483), and the Wellcome Trust (H.F., R.R., and N.P.C.).

### References


1. Tokarev, SA.; Gurchich, IS. The Yakuts. In: Levin, MG.; Potapov, LP., editors. Peoples of Siberia. University of Chicago Press; Chicago: 1956. p. 243-304.
2. Draper HH. The aboriginal Eskimo diet in modern perspective. *Am Anthropologist* 1977;79:309–16.
3. Hart TB, Hart JA. The ecological basis of hunter-gatherer subsistence in African rain forests: The Mbuti of Eastern Zaire. *Hum Ecol* 1986;14:29–55.
4. Leberthal E. Role of salivary amylase in gastric and intestinal digestion of starch. *Dig Dis Sci* 1987;32:1155–7. [PubMed: 2443325]
5. Wrangham RW, Jones JH, Laden G, Pilbeam D, Conklin-Brittain N. The raw and the stolen - cooking and the ecology of human origins. *Curr Anthropol* 1999;40:567–94. [PubMed: 10539941]
6. Klein RG. Archeology and the evolution of human behavior. *Evol Anthropol* 2000;9:17–36.
7. Diamond J. Evolution, consequences and future of plant and animal domestication. *Nature* 2002;418:700–7. [PubMed: 12167878]

8. Zohary, D.; Hopf, M. Domestication of plants in the Old World. Oxford Science Publications; Oxford: 2000.
9. Groot PC, et al. The human alpha-amylase multigene family consists of haplotypes with variable numbers of genes. *Genomics* 1989;5:29–42. [PubMed: 2788608]
10. Iafrate AJ, et al. Detection of large-scale variation in the human genome. *Nat Genet* 2004;36:949–51. [PubMed: 15286789]
11. Redon R, et al. Global variation in copy number in the human genome. *Nature* 2006;444:444–54. [PubMed: 17122850]
12. Vincent AS. Plant foods in savanna environments: a preliminary report of tubers eaten by the Hadza of northern Tanzania. *World Archaeol* 1984;17:131–148. [PubMed: 16470992]
13. Rosenberg NA, et al. Clines, clusters, and the effect of study design on the inference of human population structure. *PLoS Genet* 2005;1:e70. [PubMed: 16355252]
14. Valdez IH, Fox PC. Interactions of the salivary and gastrointestinal systems. I. The role of saliva in digestion. *Dig Dis* 1991;9:125–32. [PubMed: 1714362]
15. Read NW, et al. Swallowing food without chewing; a simple way to reduce postprandial glycaemia. *Br J Nutr* 1986;55:43–7. [PubMed: 3311145]
16. Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet* 2006;367:1747–57. [PubMed: 16731270]
17. Fried M, Abramson S, Meyer JH. Passage of salivary amylase through the stomach in humans. *Dig Dis Sci* 1987;32:1097–103. [PubMed: 3652896]
18. Samuelson LC, Wiebauer K, Snow CM, Meisler MH. Retroviral and pseudogene insertion sites reveal the lineage of human salivary and pancreatic amylase genes from a single gene during primate evolution. *Mol Cell Biol* 1990;10:2513–20. [PubMed: 1692956]
19. Cheng Z, et al. A genome-wide comparison of recent chimpanzee and human segmental duplications. *Nature* 2005;437:88–93. [PubMed: 16136132]
20. Perry GH, et al. Hotspots for copy number variation in chimpanzees and humans. *Proc Natl Acad Sci U S A* 2006;103:8006–11. [PubMed: 16702545]
21. Wilson GM, et al. Identification by full-coverage array CGH of human DNA copy number increases relative to chimpanzee and gorilla. *Genome Res* 2006;16:173–81. [PubMed: 16365383]
22. Fortna A, et al. Lineage-specific gene duplication and loss in human and great ape evolution. *PLoS Biol* 2004;2:E207. [PubMed: 15252450]
23. McGeachin RL, Akin JR. Amylase levels in the tissues and body fluids of several primate species. *Comp Biochem Physiol A* 1982;72:267–9. [PubMed: 6178549]
24. Hohmann, G.; Robbins, M.; Boesch, C. Feeding ecology in apes and other primates. Cambridge University Press; Cambridge: 2006.
25. Lambert JE. Competition, predation, and the evolutionary significance of the cercopithecine cheek pouch: the case of *Cercopithecus* and *Lophocebus*. *Am J Phys Anthropol* 2005;126:183–92. [PubMed: 15386232]
26. Coursey DG. Hominid evolution and hypogeous plant foods. *Man* 1973;8:634–5.
27. Laden G, Wrangham R. The rise of the hominids as an adaptive shift in fallback foods: plant underground storage organs (USOs) and australopithecine origins. *J Hum Evol* 2005;49:482–98. [PubMed: 16085279]
28. Hawkes K, O'Connell JF, Jones NG, Alvarez H, Charnov EL. Grandmothering, menopause, and the evolution of human life histories. *Proc Natl Acad Sci U S A* 1998;95:1336–9. [PubMed: 9448332]
29. Stranger BE, et al. Relative impact of nucleotide and copy number variation on gene expression phenotypes. *Science* 2007;315:848–53. [PubMed: 17289997]
30. Fiegler H, et al. Accurate and reliable high-throughput detection of copy number variation in the human genome. *Genome Res* 2006;16:1566–74. [PubMed: 17122085]




**Figure 1.** *AMY1* copy number variation and salivary amylase protein expression. **(a,b)** From the same European-American individuals we estimated diploid *AMY1* gene copy number with qPCR **(a)** and amylase protein levels in saliva by western blot **(b)**. Error bars indicate s.d. **(c)** Relationship between *AMY1* diploid copy number and salivary amylase protein level (n = 50 European-Americans). A considerable amount of variation in *AMY1* protein level is not explained by copy number ( $R^2 = 0.351$ ), which may reflect other genetic influences on *AMY1* expression such as regulatory region single nucleotide polymorphisms (SNPs) or non-genetic factors that may include individual hydration status, stress level, and short-term dietary habits.






**Figure 2.** Diet and *AMY1* copy number variation. **(a)** Comparison of qPCR-estimated *AMY1* diploid copy number frequency distributions for populations with traditional diets that incorporate many starch-rich foods (high-starch) and populations with traditional diets that include little or no starch (low-starch). **(b)** Cumulative distribution plot of diploid *AMY1* copy number for each of the seven populations in the study.



**Figure 3.** High-resolution fiber FISH validation of *AMY1* copy number estimates. Red (~10 kb) and green (~8 kb) probes encompass the entire *AMY1* gene and a retrotransposon directly upstream of (and unique to) *AMY1*, respectively. **(a)** Japanese individual GM18972 was estimated by qPCR to have 14 ( $13.73 \pm 0.93$ ) diploid *AMY1* gene copies, consistent with fiber FISH results showing one allele with 10 copies and the other with four copies. **(b)** Biaka individual GM10472 was estimated by qPCR to have 6 ( $6.11 \pm 0.17$ ) diploid *AMY1* gene copies, consistent with fiber FISH results. **(c)** The chimpanzee reference individual (Clint; S006006) was confirmed to have two diploid *AMY1* gene copies.



**Figure 4.** Japanese-Yakut copy number differentiation at *AMY1* versus other genome-wide loci. **(a)** Frequency distributions of WGTP aCGH relative intensity log<sub>2</sub> ratios from *AMY1*-mapped clone Chr1tp-6D2 for Japanese and Yakut individuals. **(b)** Relationship between Japanese and Yakut mean log<sub>2</sub> ratios for all autosomal WGTP clones that were copy number variable in both populations. *AMY1*-mapped clones Chr1tp-6D2 and Chr1tp-30C7 are depicted as solid red and blue circles, respectively.